

10/551,772

STN

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSPTASXK1621

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1		Web Page URLs for STN Seminar Schedule - N. America
NEWS 2		"Ask CAS" for self-help around the clock
NEWS 3	OCT 23	The Derwent World Patents Index suite of databases on STN has been enhanced and reloaded
NEWS 4	OCT 30	CHEMLIST enhanced with new search and display field
NEWS 5	NOV 03	JAPIO enhanced with IPC 8 features and functionality
NEWS 6	NOV 10	CA/CAPLUS F-Term thesaurus enhanced
NEWS 7	NOV 10	STN Express with Discover! free maintenance release Version 8.01c now available
NEWS 8	NOV 20	CA/CAPLUS to MARPAT accession number crossover limit increased to 50,000
NEWS 9	DEC 01	CAS REGISTRY updated with new ambiguity codes
NEWS 10	DEC 11	CAS REGISTRY chemical nomenclature enhanced
NEWS 11	DEC 14	WPIDS/WPINDEX/WPIX manual codes updated
NEWS 12	DEC 14	GBFULL and FRFULL enhanced with IPC 8 features and functionality
NEWS 13	DEC 18	CA/CAPLUS pre-1967 chemical substance index entries enhanced with preparation role
NEWS 14	DEC 18	CA/CAPLUS patent kind codes updated
NEWS 15	DEC 18	MARPAT to CA/CAPLUS accession number crossover limit increased to 50,000
NEWS 16	DEC 18	MEDLINE updated in preparation for 2007 reload
NEWS 17	DEC 27	CA/CAPLUS enhanced with more pre-1907 records
NEWS 18	JAN 08	CHEMLIST enhanced with New Zealand Inventory of Chemicals
NEWS 19	JAN 16	CA/CAPLUS Company Name Thesaurus enhanced and reloaded
NEWS 20	JAN 16	IPC version 2007.01 thesaurus available on STN
NEWS 21	JAN 16	WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data
NEWS 22	JAN 22	CA/CAPLUS updated with revised CAS roles
NEWS 23	JAN 22	CA/CAPLUS enhanced with patent applications from India
NEWS 24	JAN 29	PHAR reloaded with new search and display fields
NEWS 25	JAN 29	CAS Registry Number crossover limit increased to 300,000 in multiple databases
NEWS EXPRESS	NOVEMBER 10	CURRENT WINDOWS VERSION IS V8.01c, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.
NEWS HOURS		STN Operating Hours Plus Help Desk Availability
NEWS LOGIN		Welcome Banner and News Items
NEWS IPC8		For general information regarding STN implementation of IPC 8
NEWS X25		X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation

of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 16:16:32 ON 30 JAN 2007

=> file registry

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 16:16:43 ON 30 JAN 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 29 JAN 2007 HIGHEST RN 918776-45-1

DICTIONARY FILE UPDATES: 29 JAN 2007 HIGHEST RN 918776-45-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 30, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>


Uploading C:\Program Files\Stnexp\Queries\10_551772_2.str

L1 STRUCTURE UPLOADED

=> d 11

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1 sss sam

SAMPLE SEARCH INITIATED 16:17:21 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 29 TO ITERATE

100.0% PROCESSED 29 ITERATIONS

4 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**

PROJECTED ITERATIONS: 257 TO 903

PROJECTED ANSWERS: 4 TO 200

L2 4 SEA SSS SAM L1


=> d scan

L2 4 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN

IN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8aS)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-8-oxo-4-azulenyl ester (9CI)

MF C23 H30 O6


Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):2

L2 4 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
 IN Benzoic acid, 4-hydroxy-3-methoxy-, 1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI)
 MF C23 H32 O5



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L2 4 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
 IN Benzoic acid, 4-hydroxy-, 1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-1-[(2-methyl-1-oxo-2-butenyl)oxy]-4-azulenyl ester (9CI)
 MF C27 H36 O6




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):3

L2 4 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
 IN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-1-oxo-4-azulenyl ester (9CI)
 MF C22 H28 O5

Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

ALL ANSWERS HAVE BEEN SCANNED

=> s 11 sss full
 FULL SEARCH INITIATED 16:18:04 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 450 TO ITERATE

100.0% PROCESSED 450 ITERATIONS
 SEARCH TIME: 00.00.01

29 ANSWERS

L3 29 SEA SSS FUL L1

=> file caplus
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
172.55	172.76

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 16:18:12 ON 30 JAN 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 30 Jan 2007 VOL 146 ISS 6
FILE LAST UPDATED: 29 Jan 2007 (20070129/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s l3/prep

78 L3
4352731 PREP/RL
L4 15 L3/PREP
(L3 (L) PREP/RL)

=> d ibib abs hitstr 1-

YOU HAVE REQUESTED DATA FROM 15 ANSWERS - CONTINUE? Y/(N):y

L4 ANSWER 1 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:635524 CAPLUS

DOCUMENT NUMBER: 146:19979

TITLE: Ferutinin stimulates nitric oxide synthase activity in median eminence of the rat

AUTHOR(S): Colman-Saizarbitoria, Trina; Boutros, Paulo; Amesty, Angel; Bahsas, Ali; Mathison, Yaira; Garrido, Maria del Rosario; Israel, Anita

CORPORATE SOURCE: Laboratory of Bioassays and Natural Products, Laboratory of Molecular Modeling, School of Pharmacy, Universidad Central de Venezuela, Caracas, Venez.

SOURCE: Journal of Ethnopharmacology (2006), 106(3), 327-332
CODEN: JOETD7; ISSN: 0378-8741

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Several species of Ferula genus have been used in folk medicine in digestive disorders, rheumatism, headache, arthritis, and as tranquilizers, antispasmodic and aphrodisiac. From the dry and powdered roots of Ferula hermonis Boiss was extracted the oxygenated sesquiterpene 1,5-trans-daucane type: ferutinine (1). The structure of (1) was established by spectroscopic methods as: IR, 1H RMN, 13C RMN, COSY, HMBC, HMQC, NOESY, EIMS, and CIMS. The possible signaling pathway of ferutinin

(1) in nervous tissue in vitro was assessed and the results showed that this compound is able to increase nitric oxide synthase activity and inositol monophosphate accumulation (49%, each) in the median eminence of the rat brain, suggesting that compound (1) is associated to the activation of phosphoinositide breakdown and nitric oxide production (NO), the last is a gaseous intercellular messenger known to play a broad role in human biol. from homeostasis to pathol.

IT 41743-44-6P, Ferutinin


RL: NPO (Natural product occurrence); PAC (Pharmacological activity); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); USES (Uses)

(ferutinin extracted from *Ferula hermonis* increased nitric oxide synthase activity, phosphoinositide breakdown and inositol monophosphate accumulation in median eminence rat brain)

RN 41743-44-6 CAPLUS

CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:681613 CAPLUS

DOCUMENT NUMBER: 143:302470

TITLE: Rare trisubstituted sesquiterpenes daucanes from the wild *Daucus carota*

AUTHOR(S): Ahmed, Ahmed A.; Bishr, Mohktar M.; El-Shanawany, Mohamed A.; Attia, Eman Z.; Ross, Samir A.; Pare, Paul W.

CORPORATE SOURCE: Department of Chemistry, Faculty of Science, El-Minia University, El-Minia, 61519, Egypt

SOURCE: Phytochemistry (Elsevier) (2005), 66(14), 1680-1684
CODEN: PYTCAS; ISSN: 0031-9422

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Phytochem. and biol. investigation of the roots of the wild *Daucus carota* ssp. *carota* afforded three new and four known compds., including four sesquiterpenes daucane esters (1-3 [new], and 4), one polyacetylene (5), one sesquiterpene coumarin (6), and sitosterol glucoside. The structures of the new compds. were determined by comprehensive NMR studies, including DEPT, COSY, NOESY, HMQC and HMBC analyses. Based on an agar diffusion assay, 1, 2 and 4-6 were screened and found to contain a range of low antibacterial activities against four gram pos. (*Staphylococcus aureus*,

Streptomyces scabies, Bacillus subtilis, Bacillus cereus) and two gram neg. species (Pseudomonas aeruginosa, Escherichia coli) as well as antifungal against Fusarium oxysporum and Aspergillus niger using cup agar diffusion assay.


IT 864966-99-4P

RL: BSU (Biological study, unclassified); NPO (Natural product occurrence); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
(trisubstituted sesquiterpenes Daucus carota)

RN 864966-99-4 CAPLUS

CN Benzoic acid, 4-hydroxy-, (1R,3S,3aR,4R,8R,8aS)-1-(acetyloxy)-8-(benzoyloxy)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester, rel-(+)- (9CI) (CA INDEX NAME)

Rotation (+). Absolute stereochemistry unknown.
Currently available stereo shown.



REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:857419 CAPLUS

DOCUMENT NUMBER: 141:337637

TITLE: A process for the preparation of ferutinin from Ferula genus plants

INVENTOR(S): Bombardelli, Ezio; Fontana, Gabriele; Cristoni, Aldo; Mercalli, Enrico

PATENT ASSIGNEE(S): Indena S.P.A., Italy

SOURCE: PCT Int. Appl., 20 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004087179	A1	20041014	WO 2004-EP3055	20040323
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE,				

ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,
TD, TG

AU 2004226853	A1	20041014	AU 2004-226853	20040323
CA 2521100	A1	20041014	CA 2004-2521100	20040323
EP 1615651	A1	20060118	EP 2004-722572	20040323
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK				
BR 2004009166	A	20060411	BR 2004-9166	20040323
CN 1767841	A	20060503	CN 2004-80009111	20040323
JP 2006522043	T	20060928	JP 2006-504815	20040323
NO 2005004545	A	20051006	NO 2005-4545	20051003
US 2006275246	A1	20061207	US 2005-551772	20051003

PRIORITY APPLN. INFO.: IT 2003-MI661 A 20030404
WO 2004-EP3055 W 20040323

AB The invention relates to a process for the preparation of ferutinin from *Ferula* spp exts. comprising basic hydrolysis of the exts. and treatment with p-pivaloyloxybenzoic acid. The invention relates also to the use of the exts. and ferutinin in the cosmetic and dermatol. field.

IT 41743-44-6P, Ferutinin


RL: BPN (Biosynthetic preparation); COS (Cosmetic use); NPO (Natural product occurrence); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); USES (Uses)

(process for the preparation of ferutinin from *Ferula* genus plants)

RN 41743-44-6 CAPLUS

CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:690919 CAPLUS

DOCUMENT NUMBER: 141:366336

TITLE: Structure-Activity Relationships of the Estrogenic Sesquiterpene Ester Ferutinin. Modification of the Terpenoid Core


AUTHOR(S): Appendino, Giovanni; Spagliardi, Paola; Sterner, Olov; Milligan, Stuart

CORPORATE SOURCE: Dipartimento di Scienze Chimiche Alimentari Farmaceutiche e Farmacologiche, Universita del Piemonte Orientale, Novara, 28100, Italy

SOURCE: Journal of Natural Products (2004), 67(9), 1557-1564


PUBLISHER:
DOCUMENT TYPE:
LANGUAGE:
OTHER SOURCE(S):
GI

CODEN: JNPRDF; ISSN: 0163-3864
American Chemical Society
Journal
English
CASREACT 141:366336




- AB Esterification of p-hydroxybenzoic acid, a very weak estrogenic compound, with the daucane alc. jaeschkeanadiol (I; R = H) leads to a spectacular magnification of the estrogenic activity. To identify the structural elements responsible for this effect, the terpenoid core of jaeschkeanadiol p-hydroxybenzoate (ferutinin, I; R = COC6H4OH-4) was modified, capitalizing on the presence of two functionalities, the monoacylated, hydrogen-bonded 1,3-diol system and the double bond. The hydrogen bonding, while possibly useful, was not critical for activity, while hydrogenation and cyclopropanation of the double bond were tolerated. Conversely, oxidative modifications of the double bond that placed a hydroxyl on the α -face of the mol. proved detrimental. Taken together, these observations identified the substitution at C-8/C-9 as critical for activity.
- IT 109517-73-9P, 14-Hydroxyferutinin 302342-52-5P,
14-Oxoferutinin
RL: BSU (Biological study, unclassified); SPN (Synthetic preparation);
BIOL (Biological study); PREP (Preparation)
(preparation and estrogenic activity of ferutinin analogs)
- RN 109517-73-9 CAPLUS
- CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6-(hydroxymethyl)-8a-methyl-3-(1-methylethyl)-4-azulenyl ester, rel-(+)- (9CI) (CA INDEX NAME)

Rotation (+). Absolute stereochemistry unknown.




- RN 302342-52-5 CAPLUS
- CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-6-formyl-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-8a-methyl-3-(1-methylethyl)-4-azulenyl ester (9CI)
(CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2004:541439 CAPLUS
 DOCUMENT NUMBER: 141:346209
 TITLE: Structure of Samferine
 AUTHOR(S): Eshbakova, K. A.; Saidkhodzhaev, A. I.
 CORPORATE SOURCE: S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Uzbekistan
 SOURCE: Chemistry of Natural Compounds (Translation of Khimiya Prirodnikh Soedinenii) (2004), 40(2), 194-195
 CODEN: CHNCA8; ISSN: 0009-3130
 PUBLISHER: Kluwer Academic/Consultants Bureau
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



I



II

AB The authors have isolated a new compound samferine (I), of formula C₂₂H₃₀O₅, from the roots of *F. samarcandica* Korov. Alkaline hydrolysis of I produced a sesquiterpene alc. samferol (II), of the formula C₁₅H₂₆O₃, from the neutral part of the hydrolyzate.

IT 774577-57-0P, Samferine

RL: NPO (Natural product occurrence); PRP (Properties); PUR (Purification or recovery); RCT (Reactant); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); RACT (Reactant or reagent)

(isolation and mol. structure of sesquiterpene ester samferine from *F.*


samarcandica)

RN 774577-57-0 CAPLUS

CN Benzoic acid, 4-hydroxy-, (1R,3S,3aR,4R)-1,2,3,3a,4,5,8,8a-octahydro-1,3-dihydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester, rel- (9CI)
(CA INDEX NAME)

Relative stereochemistry.

Currently available stereo shown.



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:709827 CAPLUS

DOCUMENT NUMBER: 137:363261

TITLE: Daucane phytoestrogens: a structure-activity study

AUTHOR(S): Appendino, Giovanni; Spagliardi, Paola; Cravotto, Giancarlo; Pocock, Victoria; Milligan, Stuart

CORPORATE SOURCE: Dipartimento di Scienze Chimiche, Alimentari, Farmaceutiche e Farmacologiche, Universita del Piemonte Orientale, Novara, 28100, Italy

SOURCE: Journal of Natural Products (2002), 65(11), 1612-1615
CODEN: JNPRDF; ISSN: 0163-3864

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The estrogenic activity of a series of analogs of the daucane ester ferutinin modified at the acyl moiety was investigated in a yeast screen containing the human estrogen receptor α . Rather strict structure-activity relationships were observed. Thus, while the parent polyol (jaeschkeanadiol) was inactive, the presence of a p-hydroxybenzoyl moiety was necessary for activity in the yeast screen. Homologation and vinylation were both detrimental for activity, as were methylation of the p-hydroxyl substituent and the introduction of oxygen functions on the adjacent carbons.


IT 41743-44-6DP, Ferutinin, derivs. 41743-44-6P, Ferutinin
54526-95-3P 126783-56-0P

RL: PAC (Pharmacological activity); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(daucane phytoestrogens: a structure-activity study)

RN 41743-44-6 CAPLUS

CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 41743-44-6 CAPLUS

CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 54526-95-3 CAPLUS


CN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 126783-56-0 CAPLUS
CN Benzoic acid, 3,4-dihydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 7 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2002:315620 CAPLUS
DOCUMENT NUMBER: 137:332761
TITLE: Antimicrobial sesquiterpene from the roots of *Ferula harmonis*
AUTHOR(S): Al-Sha'er, M.; Darwish, R. M.; Aburjai, T.
CORPORATE SOURCE: Department of Pharmaceutical Sciences, Faculty of Pharmacy, University of Jordan, Amman, 11942, Jordan
SOURCE: Acta Technologiae et Legis Medicamenti (2001), 12(3), 255-264
CODEN: ATLMEQ; ISSN: 1121-2098
PUBLISHER: Maccari Editore
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Two known sesquiterpenes were isolated for the first time from the roots of *Ferula harmonis* F. (Umbelliferae). Their structures were assigned by spectroscopic (MS, ¹H-NMR, ¹³C-NMR) and phys. means as ferutin and teferidine. The antimicrobial activity of the crude exts. and the isolated compds. was tested against four different microorganisms. Ferutin showed good antibacterial activity against *S. aureus* and weak antifungal activity against *A. niger*. Teferidine, showed no antimicrobial activity against the tested microorganisms. Our results revealed the importance of the phenolic hydroxyl group for the antimicrobial activity of these compds.

IT 41743-44-6P, Ferutin
RL: NPO (Natural product occurrence); PAC (Pharmacological activity); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); USES (Uses)
(antimicrobial sesquiterpene from the roots of *Ferula harmonis*)

RN 41743-44-6 CAPLUS
CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 8 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2001:771413 CAPLUS

DOCUMENT NUMBER: 136:180636

TITLE: Sesquiterpenoids from the fruits of *Ferula kuhistanica* and antibacterial activity of the constituents of *F. kuhistanica*

AUTHOR(S): Tamemoto, Kimiko; Takaishi, Yoshihisa; Chen, Bei; Kawazoe, Kazuyoshi; Shibata, Hirofumi; Higuti, Tomihiko; Honda, Gisho; Ito, Michiho; Takeda, Yoshio; Kodzhimatov, Olimjon K.; Ashurmetov, Ozodbek

CORPORATE SOURCE: Faculty of Pharmaceutical Sciences, University of Tokushima, Tokushima, 770-8505, Japan

SOURCE: Phytochemistry (2001), 58(5), 763-767

CODEN: PYTCAS; ISSN: 0031-9422

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Et acetate exts. of the air-dried fruits of *Ferula kuhistanica* afforded three daucane esters: kuhistanicaols H, I and J (I, II, and III, resp.), together with nine other known compds. Their structures were established on the basis of spectroscopic evidence. Isolated compds. in this paper and previously reported compds. from the roots and stems of *F. kuhistanica* were tested for antibacterial activity. Some of them were selectively toxic against Gram-pos. bacteria, including methicillin-sensitive and methicillin-resistant *Staphylococcus aureus* (MSSA and MRSA).

IT 41743-44-6P


RL: PAC (Pharmacological activity); PRP (Properties); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(purification from *Ferula kuhistanica* fruit and antibacterial activity of)

RN 41743-44-6 CAPLUS


CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 54526-95-3P
 RL: PAC (Pharmacological activity); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation);
 USES (Uses)
 (purification from *Ferula kuhistanica* fruit and antibacterial activity of)
 RN 54526-95-3 CAPLUS
 CN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.



IT 399035-10-0P, Kuhistanicaol I
 RL: NPO (Natural product occurrence); PAC (Pharmacological activity); PRP (Properties); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); USES (Uses)
 (sesquiterpenoids from the fruit of *Ferula kuhistanica* and antibacterial activity of the constituents of *F. kuhistanica*)
 RN 399035-10-0 CAPLUS
 CN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8R,8aS)-1,2,3,3a,4,5,8,8a-octahydro-3,8-dihydroxy-6,8a-dimethyl-3-(1-methylethyl)-1-oxo-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 9 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2001:286907 CAPLUS

DOCUMENT NUMBER: 135:43518

TITLE: Daucane aryl esters composition from the Lebanese *Ferula hermonis* Boiss. (zallooh root)

AUTHOR(S): Diab, Youssef; Dolmazon, Rene; Bessiere, Jean-Marie

CORPORATE SOURCE: Universite Libanaise, Faculte des Sciences-2, Jdaidet el-Matn, Lebanon

SOURCE: Flavour and Fragrance Journal (2001), 16(2), 120-122
CODEN: FFJOED; ISSN: 0882-5734

PUBLISHER: John Wiley & Sons Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The dichloromethane extract of *Ferula hermonis* roots, isolated in 26.5% yield based on dry roots, contained as principal components p-hydroxybenzoate and benzoate of jaeschkeanadiol (52% and 30%, resp.). Some other aryl esters of jaeschkeanadiol and 2,3-epoxide jaeschkeanadiol, as well as α -bisabolol, were present in small amts.

IT 41743-44-6P, Jaeschkeanadiol p-hydroxybenzoate 54526-95-3P
, Teferin


RL: BOC (Biological occurrence); BSU (Biological study, unclassified); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)

(daucane aryl esters composition from the Lebanese *Ferula hermonis* (zallooh root))

RN 41743-44-6 CAPLUS


CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 54526-95-3 CAPLUS
CN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 10 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2001:4915 CAPLUS
DOCUMENT NUMBER: 134:219768
TITLE: Sesquiterpenes from *Ferula hermonis* Boiss
AUTHOR(S): Galal, A.
CORPORATE SOURCE: National Center for Natural Products Research, School of Pharmacy, University of Mississippi, University, MS, 38677, USA
SOURCE: Pharmazie (2000), 55(12), 961-962
CODEN: PHARAT; ISSN: 0031-7144
PUBLISHER: Govi-Verlag Pharmazeutischer Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English
AB The roots of *Ferula hermonis* yielded the new 8,9-epoxy derivative of the carotane sesquiterpene jaeschkeanadiol, together with 2 other known sesquiterpenes: (+)- α -bisabolol and jaeschkeanadiol vanillate. The identities of the isolated compds. were established from their spectral data and by comparison with published reports.
IT 54526-95-3P
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
(sesquiterpenes from *Ferula hermonis*)
RN 54526-95-3 CAPLUS
CN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 11 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:590836 CAPLUS
 DOCUMENT NUMBER: 133:319520
 TITLE: Sesquiterpenoids from *Ferula kuhistanica*
 AUTHOR(S): Chen, B.; Teranishi, R.; Kawazoe, K.; Takaishi, Y.;
 Honda, G.; Itoh, M.; Takeda, Y.; Kodzhimatov, O. K.
 CORPORATE SOURCE: Fac. Pharm. Sci., Univ. Tokushima, Tokushima,
 770-8505, Japan
 SOURCE: Phytochemistry (2000), 54(7), 717-722
 CODEN: PYTCAS; ISSN: 0031-9422
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



I

AB Methanol exts. of the air-dried roots and stems of *Ferula kuhistanica* afforded seven daucane-type sesquiterpenes, called kuhistanicaol A-G (e.g. I, kuhistanicaol A) together with 13 known daucane esters. Their structures were established on the basis of spectroscopic evidence and the results of chemical reactions.
 IT 302342-52-5P, Kuhistanicaol B 302342-54-7P,
 Kuhistanicaol D
 RL: BOC (Biological occurrence); BSU (Biological study, unclassified); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
 (from *Ferula kuhistanica*)
 RN 302342-52-5 CAPLUS
 CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-6-formyl-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-8a-methyl-3-(1-methylethyl)-4-azulenyl ester (9CI)
 (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 302342-54-7 CAPLUS

CN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8S,8aS)-1,2,3,3a,4,5,8,8a-octahydro-3,8-dihydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 12 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:294524 CAPLUS
Correction of: 1998:104683

DOCUMENT NUMBER: 130:294014
Correction of: 128:203022

TITLE: Chemical constituents of rocks of *Ferula licentiana* var. *tunshanica* and *F. kingdonwardii* and their systematical significance

AUTHOR(S): Wang, Nianhe; Yuan, Changqi; Baba, Kimie; Taniguchi, Masahiko; Doi, Mitsunobu

CORPORATE SOURCE: Institute of Botany, Jiangsu Province and Chinese Academy of Sciences, Nanjing, 210014, Peop. Rep. China

SOURCE: *Zhiwu Ziyuan Yu Huanjing* (1997), 6(4), 15-18, 49
CODEN: ZZYEJ; ISSN: 1004-0978

PUBLISHER: Zhiwu Ziyuan Yu Huanjing Bianjibu

DOCUMENT TYPE: Journal

LANGUAGE: Chinese

AB Latifolone, p-anisic acid, veratric acid, fercomin, ferutidin, ferutinin, jaeschkeanadiol varatrate, terferin, 1-(7,8-dimethoxy-5,6-methylenedioxy phenyl)propyl (Z)-2-methyl-1-butenate and 4,11,11,10-tetramethyl-1,10-oxirane-4-ene-6-germacrane varatrate were isolated from the ether exts. of the roots of *Ferula licentiana* Hand.-Mazz. var. *tunshanica* Shan et Q. X. Liu and *F. kingdonwardii* Wolff. Among them, 1-(7,8-dimethoxy-5,6-

methylenedioxy phenyl) Pr (Z)-2-methyl-butenolate is a new compound. These two Ferulic plants were distributed on the border of the distribution of this genus, but 7-O-sesquiterpene, one of the characteristic compounds of this genus, was not detectable in these species, and the components of these two plants were similar and comparatively simple. It was suggested that perhaps *Ferula* L., like some other Umbelliferous plants such as *Angelica* L., also originated from the south-west China.

IT 41743-44-6P, Ferutin 54526-95-3P, Teferin


RL: BOC (Biological occurrence); BSU (Biological study, unclassified); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)

(chemical constituents of roots of *ferula licentiana* var. *tunshanica* and *F. kingdonwardii* and systematical significance)

RN 41743-44-6 CAPLUS

CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 54526-95-3 CAPLUS

CN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




ACCESSION NUMBER: 1998:104683 CAPLUS
 DOCUMENT NUMBER: 128:203022
 TITLE: Chemical constituents of roots of *Ferula licentiana* var. *tunshanica* and *F. kingdonwardii* and their systematical significance
 AUTHOR(S): Wang, Nianhe; Yuan, Changqi; Kimie Baba; Masahiko Taniguchi; Mitsunobu Doi
 CORPORATE SOURCE: Institute of Botany, Jiangsu Province and Chinese Academy of Sciences, Nanjing, 210014, Peop. Rep. China
 SOURCE: Zhiwu Ziyuan Yu Huanjing (1997), 6(4), 15-18, 49
 CODEN: ZZYEJ; ISSN: 1004-0978
 PUBLISHER: Zhiwu Ziyuan Yu Huanjing Bianjibu
 DOCUMENT TYPE: Journal
 LANGUAGE: Chinese

AB Latifolone, p-anisic acid, veratric acid, fercomin, ferutidin, ferutinin, jaeschkeanadiol varatrate, terferin, 1-(7,8-dimethoxy-5,6-methylenedioxy phenyl)propyl (Z)-2-methyl-1-butenolate and 4,11,11,10-tetramethyl-1,10-oxirane-4-ene-6-germacrane varatrate were isolated from the ether exts. of the roots of *Ferula licentiana* Hand.-Mazz. var. *tunshanica* Shan et Q. X. Liu and *F. kingdonwardii* Wolff. Among them, 1-(7,8-dimethoxy-5,6-methylenedioxy phenyl) Pr (Z)-2-methyl-butenolate is a new compound. These two *Ferulious* plants were distributed on the border of the distribution of this genus, but 7-O-sesquiterpene, one of the characteristic compds. of this genus, was not detectable in these species, and the components of these two plants were similar and comparatively simple. It was suggested that perhaps *Ferula* L., like some other Umbelliferous plants such as *Angelica* L., also originated from the south-west China.

IT 41743-44-6P, Ferutinin 54526-95-3P, Teferin
 RL: BOC (Biological occurrence); BSU (Biological study, unclassified); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
 (chemical constituents of roots of *ferula licentiana* var. *tunshanica* and *F. kingdonwardii* and systematical significance)


RN 41743-44-6 CAPLUS
 CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 54526-95-3 CAPLUS
 CN Benzoic acid, 4-hydroxy-3-methoxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 14 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1995:844841 CAPLUS

DOCUMENT NUMBER: 124:4913

TITLE: Carotane sesquiterpenes from *Ferula sinaica* Boiss growing in Egypt

AUTHOR(S): Ibraheim, Zidan Z.; Darwish, Faten M. M.; Abdel-Halim, Osama B.; Halim, Ahmed F.

CORPORATE SOURCE: Faculty Pharmacy, Assiut University, Assiut, Egypt

SOURCE: Alexandria Journal of Pharmaceutical Sciences (1995), 9(2), 115-20

CODEN: AJPSES; ISSN: 1110-1792

PUBLISHER: University of Alexandria, Faculty of Pharmacy

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Phytochem. investigation of the air-dried roots of *Ferula sinaica* Boiss resulted in the isolation and identification of ten known carotane sesquiterpenes. Of these, seven compds. were isolated for the first time from this species and characterized as: 1- α -angeloyloxy-5- α -p-methoxybenzoyloxy-10- β -hydroxydauc-2-ene, lancerodiol p-methoxybenzoate, p-methoxybenzoate of epoxyjaeschkeanadiol, p-hydroxybenzoate of epoxyjaeschkeanadiol, isolancerotriol-5-p-methoxybenzoate, 1 α - β -hydroxy-5- α -p-methoxybenzoyloxy-10- β -hydroxydauc-2-ene and 1- α -hydroxy-5- α -p-methoxybenzoyloxy-10- β -hydroxydauc-2-ene in addition to β -sitosterol. Identification of the isolated compds. was established using phys. and spectroscopic methods.


IT 41743-44-6P 119425-93-3P 170971-43-4P,
Ferulinkiol 1- α -hydroxy-5- α -p-hydroxybenzoate
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)

(carotane sesquiterpenes from *Ferula sinaica* growing in Egypt)

RN 41743-44-6 CAPLUS


CN Benzoic acid, 4-hydroxy-, (3R,3aS,4S,8aR)-1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 119425-93-3 CAPLUS
 CN Benzoic acid, 4-hydroxy-, 1,2,3,3a,4,5,8,8a-octahydro-3-hydroxy-6,8a-dimethyl-3-(1-methylethyl)-8-[(2-methyl-1-oxo-2-butenyl)oxy]-4-azulenyl ester, [3R-[3 α ,3 β ,4 β ,8 β (Z),8 α]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.




RN 170971-43-4 CAPLUS
 CN Benzoic acid, 4-hydroxy-, 1,2,3,3a,4,5,8,8a-octahydro-3,8-dihydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester, (3 α ,3 β ,4 β ,8. beta.,8 α)- (9CI) (CA INDEX NAME)

Relative stereochemistry.
 Currently available stereo shown.




L4 ANSWER 15 OF 15 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1980:471992 CAPLUS
 DOCUMENT NUMBER: 93:71992
 TITLE: Structure of jaeschferin
 AUTHOR(S): Bizhanova, K. B.; Saidkhodzhaev, A. I.; Malikov, V. M.
 CORPORATE SOURCE: Inst. Khim. Rastit. Veshchestv, Tashkent, USSR
 SOURCE: Khimiya Prirodnikh Soedinenii (1980), (1), 127-8
 CODEN: KPSUAR; ISSN: 0023-1150
 DOCUMENT TYPE: Journal
 LANGUAGE: Russian
 GI



AB The structure of jaeschferin (I), a sesquiterpene ester isolated from
 Ferula iaeschkeana, was confirmed by mass and NMR spectra.
 IT 74345-97-4P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 74345-97-4 CAPLUS
 CN Benzoic acid, 4-hydroxy-3-methoxy-, 1,2,3,3a,4,5,8,8a-octahydro-1,3-
 dihydroxy-6,8a-dimethyl-3-(1-methylethyl)-4-azulenyl ester, diacetate,
 (1 α ,3 α ,3a β ,4 β ,8a α) - (9CI) (CA INDEX NAME)
 CM 1
 CRN 74285-96-4
 CMF C23 H32 O6


Rotation (+). Absolute stereochemistry unknown.



CM 2

CRN 64-19-7

CMF C2 H4 O2



=> d his

(FILE 'HOME' ENTERED AT 16:16:32 ON 30 JAN 2007)

FILE 'REGISTRY' ENTERED AT 16:16:43 ON 30 JAN 2007

L1 STRUCTURE UPLOADED

L2 4 S L1 SSS SAM

L3 29 S L1 SSS FULL

FILE 'CAPLUS' ENTERED AT 16:18:12 ON 30 JAN 2007

L4 15 S L3/PREP

=>